### **Health Protection Hub**

# Guide for Schools, Colleges and Nurseries Dealing with Diarrhoea & Vomiting

#### Introduction

This guide is designed to be a single easy-to-use resource for staff in educational settings dealing with an outbreak of Diarrhoea and Vomiting. Usually these are viral in nature, such as Norovirus, but may be linked to consumption of food or water and require some investigation and control measures to reduce transmission.

### Steps to take

- 1. Advise the HPH by calling 01952 381800 or emailing <a href="mailto:HealthProtectionHub@telford.gov.uk">HealthProtectionHub@telford.gov.uk</a>
- 2. Work through the checklist at Appendix 1 (which may require you to liaise with kitchen and/or cleaning contractors), email it in to the two addresses at the top of the document and address any areas of shortfall identified by the process.
- 3. Draft a simple analysis for the affected cases and send in to the Health Protection Hub example template below. Even if the information is incomplete this will aid in an understanding of the situation.

Initials	School	Class/room	Symptoms	Symptom	Date last in
	dinners (Y/N)			onset date	setting
ST	Y	4	Vomiting	8/2	8/2
BY	N	4	Vomiting	9/2	9/2
RY	N	2	Diarrhoea	10/2	9/2
TR	N	4	Vomiting	10/2	10/2
PB	Y	4	Diarrhoea	10/2	9/2

- 4. Check that no kitchen staff have any symptoms (including boils, sores or open wounds) and exclude them from work if necessary.
- 5. Send the letter in Appendix 2 below out to parents.
- 6. If you contract-out your cleaning, advise the supervisor so they are aware and can act accordingly. They may be able to arrange additional cleaning such as steam cleaning of soft furnishings if necessary.
- 7. Contact the parent/carers of children who are off from school or nursery without a reason and find out if they have symptoms.
- 8. Attempt to rule out COVID-19 by asking parents to test using home LFD kits and report back the results for the purposes of excluding it as the source of the infection (this is covered in the template letter above).
- 9. Keep the table in Point 3 above updated with new cases.
- 10. Depending on the situation you may be visited by Environmental Health Officers to check food hygiene measures in the kitchen and/or the wider measures in place to control the outbreak. You may also be asked for contact details of affected people so that stool samples can be obtained for analysis.
- 11. If the case numbers don't drop within a couple of days, or you want any further advice at any point, feel free to contact the Health Protection Hub.

# Appendix 1

Checklist completed by		
Name/position:		
Date:		
Setting: Nursery/School - Primary/		
Secondary/ SEN or FE College		
Local Authority/Private/Academy		
If special school, nature of special needs.		
Name, address and postcode		
Confined to one class year group or throughout setting?	Yes / No	
Pupils confirmed or suspected out	No. of pupils confirmed	
of total attending setting/ year	Total No. of pupils in	
group affected –	setting	
Staff confirmed or suspected out of	No. of staff confirmed	
total employed	Tatal Na of staff in	
	Total No. of staff in	
Opent data of superstance in the first	setting	
Onset date of symptoms in the first		
case		
Is the first case a member of staff or pupil?		
Symptoms: Vomiting Diarrhoea		
Fever other		
Any common foods consumed		
within setting in previous 48		
hours? If yes list foods below		
Common Foods		
Have cases have tested positive for		
COVID-		

Potential cohorts affected: e.g.	Year Group/	Total no. of pupils
<ul> <li>class/year group</li> </ul>	Class	attending
<ul> <li>breakfast or after-school clubs</li> </ul>		
(extra-curricular activity)		
<ul> <li>school transport</li> </ul>		
<ul> <li>Do any of the cases attend more</li> </ul>		
than one setting?		
School trips		
Have any cases submitted a faecal		
sample		
Any cases hospitalised or had a		
confirmed diagnosis		
Communication with staff and		
students to date ( template letter		
sent )		
Any media interest or concerns		
from pupils, parents or staff?		

# Please tick if action is complete and add any comments

Outbreak control:	
Parents/carers of children that are ill whilst at school should	
be contacted and requested that they come to collect their	
child from school or nursery.	
Affected children should be isolated, if possible from their	
class mates until collected by their parents/carer.	
Children and Staff that have Norovirus symptoms must	
remain off school or nursery for 48 Hours after the last	
episode of vomiting or diarrhoea	
Sections in red are required control measures	
Hand and respiratory hygiene	
Are there appropriate facilities for handwashing – hand	
sanitisers, liquid soap and paper towels/tissues and pedal	
operated bins For teaching children use e-Bug coronavirus	
(COVID-19) website	
A handwashing programme should be put into place that	
encourages children to wash their hands at the start of the	
school day, after using the toilet, after play, before and after	
eating, after touching pets and animals, and encouraging	
parents to let their children wash their hands at the end of	
the school day	

	Annual company shift damping and shift data are 1900.	
	Are younger children and children with complex needs	
	supervised, helped with hand hygiene?	
	Cleaning	
	The frequency of toilet cleaning should be increased, eg	
	clean after each break time and after a child has been sick or	
	had diarrhoea.	
	Touch points, e.g. taps, toilet flush handles, door handles,	
	should be cleaned regularly with a hypochlorite (bleach	
	based) solution 1,000 parts per million. Read manufacturer's	
	instructions and do not use on fabrics and carpets	
	Soiled children's clothing should be sealed in a plastic bag to	
	go home. It should not be washed on site	
	Potties should be cleaned and disinfected after use and	
	should only be used by the same child. If this is not possible	
	they should be cleaned and disinfected after use with 1000	
	parts per million hypochlorite (bleach-based solution)	
	Toys used by the children should be washed and if possible	
	disinfected. Soft toys should be machine washable; hard	
	surface toys are more easily washed and disinfected.	
	Advice on cleaning after a suspected/confirmed case has left	
	setting. Clean surfaces with a household detergent followed	
	by disinfection (1000 ppm available chlorine or a disinfectant	
	,	
	that works against enveloped viruses). Follow	
	manufacturer's instructions for dilution, application and	
	contact times for all detergents and disinfectants	
	Single use disposable cloths and mop heads to be used, if not	
	available cloths and mop heads to be laundered after use.	
	Spray into a cloth not onto surfaces, avoid creating splashes	
	and spray when cleaning	
	When items cannot be cleaned using detergents or	
	laundered, for example, upholstered furniture, steam	
	cleaning should be used	
	Classrooms	
	Staff movements between classrooms and joint class	
	activities in school, e.g. assembly should be restricted	
	Are classrooms clutter free? Soft furnishings, soft toys and	
	toys that are hard to clean should not be in use during an	
	outbreak.	
	To avoid if possible, sharing of resources such as toys/	
	keyboards with other groups/classes. These should be	
	cleaned between use.	
	Equipment shared between classes to be cleaned frequently	
	and always between classes/groups or rotated to allow them	
	to be left unused and out of reach for a period of 48 hours	
	(72 hours for plastics).	
<u> </u>	, ,	1

Stop sand and water play, use of play dough / plastic cookery lessons. Sand, play dough and plasticine sho thrown away	
Consider whether any activities could take place out including exercise, assemblies, or classes	doors,
Visits to the setting	
Unless essential, stop any visitors attending the sett	ng.
Group visits in and out of school should be stopped hours after the last person has stopped having symp	
Disposable aprons and gloves should be worn when touch points, cleaning and disinfecting potties, chan nappies, toileting a child or cleaning up vomit or dial Staff should wash their hands after the removal of gaprons.	ging rhoea.
Is there adequate supplies of PPE, gloves, aprons, meeye protection?	asks and
Food/drink	
Food should be either prepared by canteen staff or lin by a child and consumed by that child only, i.e. no of food	
Drinking utensils are for individual use only.	
Special precautions and supervision in filling water be ensure taps are not contaminated.	ottles to
Discourage use of water fountains for drinking water	۲.
Crockery and cutlery to be dishwasher washed and s cupboards	tored in
Keep all food in cupboards or in a lidded wipeable co do not use shared fruit bowls etc	ontainer,
Caring for a symptomatic child	
If a child develops symptoms whilst at school and is collection isolate the child in a ventilated room with supervision or move them to an area which is at leas metres away from other people.	staff It two
PPE ( gloves and aprons) should be worn by staff car the child while they await collection	ing for
Handling of waste	
Waste from possible/confirmed and cleaning of area possible cases have been (including disposable cloth heads. PPE used, and tissues used by case etc.) to be bagged and tied. Remove to designated storage area	s, mop double

### Appendix 2

Dear parent/carer

This letter is to advise you that we have cases of diarrhoea and vomiting in the setting and are liaising with the Health Protection Hub and taking measures to control it. From the information we currently have we believe that it is a viral illness such as Norovirus (otherwise known as the winter vomiting bug).

Below is some information on Norovirus and what you can do to reduce the spread of the infection:

#### What are the symptoms?

Diarrhoea and vomiting usually lasting 1-2 days

## What should you do if your child is unwell?

Make sure they get lots of rest.

Ensure they drink plenty of fluids, taking sips rather than gulps to avoid vomiting.

Give infant Paracetamol or Ibuprofen, according to product instructions, to help keep their temperature down.

If your child is unusually sleepy, won't take fluids or has other symptoms, such as blood in their diarrhoea, an unusual rash, headache, neck stiffness or difficulty breathing, ring NHS Direct (0845 4647), contact your GP or take your child to hospital.

If any siblings are unwell please contact their nursery/ school and let them know that your child is ill and what symptoms they have.

Keep them at home until they are well enough to return to school. However, if your child has been suffering from diarrhoea and vomiting keep them at home until they have been free of symptoms for 48 hours.

If you have LFD (test kits) at home test your child for COVID-19 and let the school know the result if possible so they can potentially exclude it as the cause.

### How can you prevent the spread of these infection?

Hand washing is one of the most important ways to prevent the spread of both of these infections. This applies to the child who is ill and the person looking after them. Hands should always be washed, using liquid soap if possible: Before and after caring for your child, after using the toilet, before eating or handling food, after cleaning up a mess (vomit, faeces or urine).

Other ways of preventing the spread of infection are to:

Keep a separate towel for family members who have symptoms.

Dispose promptly of used tissues and other items that your child may have coughed or sneezed on.

Wash soiled clothing, bed linen and towels in a washing machine. Use the hottest wash for that fabric.

Clean baths and washbasins thoroughly and disinfect after use.

If cleaning up diarrhoea or vomit, wash the surface with hot soapy water and disinfect before allowing to dry. If using bleach remember that it can remove the colour from fabrics and can burn the skin.

Keep your child away from other children and people that are particularly vulnerable, e.g. the elderly and those with chronic illnesses.

Yours faithfully